

REPORT DOCUMENTATION PAGE					Form Approved OMB No. 0704-0188	
<p>The public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden, to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.</p> <p>PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.</p>						
1. REPORT DATE (DD-MM-YYYY)		2. REPORT TYPE Final			3. DATES COVERED (From - To)	
4. TITLE AND SUBTITLE Nearshore Circulation in Complex Regions				5a. CONTRACT NUMBER		
				5b. GRANT NUMBER		
				5c. PROGRAM ELEMENT NUMBER 602435N		
				5d. PROJECT NUMBER		
6. AUTHOR(S) Kaihatu, James M. and Rogers, Erick				5e. TASK NUMBER 03550		
				5f. WORK UNIT NUMBER		
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Naval Research Laboratory Oceanography Division Stennis Space Center, MS 39529-5004				8. PERFORMING ORGANIZATION REPORT NUMBER NRL/PU/7320/01/0002		
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Office of Naval Research 800 N. Quincy St. Arlington, VA 22217-5660				10. SPONSOR/MONITOR'S ACRONYM(S) ONR		
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)		
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution is unlimited						
13. SUPPLEMENTARY NOTES						
<div style="text-align: right; font-size: 2em; font-weight: bold; margin-top: 20px;">20021101 018</div>						
14. ABSTRACT As ocean surface waves propagate over the continental shelf and break in shallow nearshore areas, their energy is dissipated in the surf zone. The gradients in the wavelets and momentum give rise to the generation of wave-induced nearshore circulation Wave forcing of circulation in the surf zone is a major cause of sediment transport and beach morphology evolution; its understanding and prediction are essential from military application such as amphibious landings and mine warfare. Semiempirical expressions for the nearshore circulation exist over planar bathymetry. However, over more complex bathymetric configurations, such as sandbars, rip channels, and canyons, numerical models are required form comprehensive descriptions of the wave-driven flow. Most of these extant numerical models either use irregular wave forcing but simplify the hydrodynamics (as is the cases for nearshore circulation models in the operational Navy), or allow more involved hydrodynamic formulations but reduce the forcing to that of monochromatic wave theory. Wither modeling option can potentially lead to nearshore hydrodynamic predictions that do not exhibit the variability seen in nature.						
15. SUBJECT TERMS surf zone, mine warfare, amphibious landings						
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT		18. NUMBER OF PAGES	
a. REPORT Unclassified	b. ABSTRACT Unclassified	c. THIS PAGE Unclassified	SAR		2	
					19a. NAME OF RESPONSIBLE PERSON Jim Kaihatu	
					19b. TELEPHONE NUMBER (Include area code) (228) 688-5710	

PUBLICATION OR PRESENTATION RELEASE REQUEST

Pubkey: 3077

NRLINST 5600.2

1. REFERENCES AND ENCLOSURES	2. TYPE OF PUBLICATION OR PRESENTATION	3. ADMINISTRATIVE INFORMATION
Ref: (a) NRL Instruction 5600.2 (b) NRL Instruction 5510.40D	() Abstract only, published () Book () Conference Proceedings (refereed) () Invited speaker () Journal article (refereed) () Oral Presentation, published (X) Other, explain <u>NRL Publication</u>	STRN <u>NRL/PU/7320-01-2</u> Route Sheet No. <u>7320/</u> Job Order No. _____ Classification <u>X</u> U _____ C Sponsor _____ approval obtained <u>X</u> yes _____ no
Encl: (1) Two copies of subject paper (or abstract)	() Abstract only, not published () Book chapter () Conference Proceedings (not refereed) () Multimedia report () Journal article (not refereed) () Oral Presentation, not published	

4. AUTHOR

Title of Paper or Presentation

Nearshore Circulation in Complex Regions

Author(s) Name(s) (First, MI, Last), Code, Affiliation if not NRL

James M Kaihatu, Erick Rogers

It is intended to offer this paper to the _____

(Name of Conference)

(Date, Place and Classification of Conference)

and/or for publication in 2002 NRL Review Book, Unclassified

(Name and Classification of Publication)

(Name of Publisher)

After presentation or publication, pertinent publication/presentation data will be entered in the publications data base, in accordance with reference (a).

It is the opinion of the author that the subject paper (is _____) (is not X) classified, in accordance with reference (b).This paper does not violate any disclosure of trade secrets or suggestions of outside individuals or concerns which have been communicated to the Laboratory in confidence. This paper (does _____) (does not X) contain any militarily critical technology.This subject paper (has _____) (has never X) been incorporated in an official NRL Report.

James M Kaihatu, 7322

Name and Code (Principal Author)

(Signature)

5. ROUTING/APPROVAL

CODE	SIGNATURE	DATE	COMMENTS
Author(s) Kaihatu		11/30/01	
Section Head Preller		12/5/01	
Branch Head John M. Harding, 7320		12/6/01	
Division Head William J. Jobst, 7300		12/7	1. Release of this paper is approved. 2. To the best knowledge of this Division, the subject matter of this paper (has _____) (has never <u>X</u>) been classified.
Security, Code 7030.1		12/11	1. Paper or abstract was released. 2. A copy is filed in this office. <u>SSC 416</u> - 01
Office of Counsel, Code 1008.3		12/17/01	Memorandum
ADOR/Director NCST E.O. Hartwig, 7000			
Public Affairs (Unclassified/Unlimited Only), Code 7030.4		12/17/01	acknowledgment
Division, Code			
Author, Code			



2002

NRL REVIEW

ARCH
the Navy's
corporate laboratory

NAVAL RESEARCH LABORATORY

Study of Microbial Chromium(VI) Reduction by Electron Energy Loss Spectroscopy	115
<i>T.L. Daulton, B.J. Little, and J.M. Jones-Meehan</i>	

ELECTRONICS AND ELECTROMAGNETICS

Digital Array Radar: A New Vision	121
<i>J.W. de Graaf and B.H. Cantrell</i>	
Traps in GaN-based Microwave Devices	122
<i>P.B. Klein, S.C. Binari, K. Ikossi, D.D. Koleske, A.E. Wickenden, and R.L. Henry</i>	
Polar Reformatting for ISAR Imaging	124
<i>R. Lipps and M. Bottoms</i>	

ENERGETIC PARTICLES, PLASMAS, AND BEAMS

The Electra KrF Laser Program	129
<i>J.D. Sethian, M. Myers, M. Friedman, R. Lehmberg, J. Giuliani, S. Obenschain, F. Hegeler, S. Swanekamp, and D. Weidenheimer</i>	
Charging and Shielding of "Dust Grains" in a Plasma	131
<i>M. Lampe, G. Ganguli, and G. Joyce</i>	
An Electrodeless Moly-Oxide Discharge for Lighting Applications	133
<i>J.L. Giuliani, R.A. Meger, R.E. Pechacek, and G.M. Petrov</i>	

INFORMATION TECHNOLOGY AND COMMUNICATIONS

Evaluation of Electronic Documents for Preparing Naval Meteorological and Oceanographic Briefings	139
<i>J.A. Ballas, W.C. Kooiman, and R.T. Miyamoto</i>	
Satellite Networking for Naval Battlegroups	141
<i>M.A. Rugar, A.S. Eley, and M. Solsman</i>	
Advanced Visualization for Test and Evaluation and Training Ranges	143
<i>J.Q. Binford and W.A. Doughty</i>	
Signal Sorter for Advanced Multifunction Radio Frequency Concept (AMRF-C) Using Neural Networks and Advanced Statistical Techniques	146
<i>V.C. Kowtha, M.J. Thompson, T.N. Reynolds, J.R. Connell, A.E. Spezio, and J.C. Sciortino, Jr.</i>	

MATERIALS SCIENCE AND TECHNOLOGY

Protectively Coated Phosphors for Flat Panel FED Devices	153
<i>J.S. Sanghera, G. Villalobos, S.S. Bayya, and I.D. Aggarwal</i>	
Scanning Nanomechanics	155
<i>K.J. Wahl, S.A. Syed Asif, and R.J. Colton</i>	
Raman Spectroscopy of High-Temperature Superconductors	157
<i>C. Kendziora</i>	
Quantum Dot Bioconjugates in Molecular Detection	159
<i>J.M. Mauro, G.P. Anderson, E.R. Goldman, H. Mattoussi, and B.L. Justus</i>	
Selective Resputtering-Induced Magnetic Anisotropy in High-Density Magneto-optic Media	161
<i>V.G. Harris</i>	

OCEAN SCIENCE AND TECHNOLOGY

Anatomy of the Ocean Surface Roughness	165
<i>P.A. Hwang, D.W. Wang, W.J. Teague, and G.A. Jacobs</i>	
Nearshore Circulation in Complex Regions	167
<i>J.M. Kaihatu and W.E. Rogers</i>	
Remote Wind Connections to Strait Transports	169
<i>G.A. Jacobs, H.T. Perkins, R.H. Preller, H.E. Ngodock, W.J. Teague, S.K. Reidlinger, D. Ko, and J.W. Book</i>	
Laboratory for Underwater Hydrodynamics	171
<i>J. Grun, T. Jones, C. Manka, and L.D. Bibee</i>	

NEARSHORE CIRCULATION IN COMPLEX REGIONS

J.M. Kaihatu and W.E. Rogers
Oceanography Division

Introduction: As ocean surface waves propagate over the continental shelf and break in shallow nearshore areas, their energy is dissipated in the surf zone. The gradients in the waveheights and momentum give rise to the generation of wave-induced nearshore circulation. Wave forcing of circulation in the surf zone is a major cause of sediment transport and beach morphology evolution; its understanding and prediction are essential for military applications such as amphibious landings and mine warfare. Semi-empirical expressions for the nearshore circulation exist over planar bathymetry. However, over more complex bathymetric configurations, such as sandbars, rip channels, and canyons, numerical models are required for comprehensive descriptions of the wave-driven flow. Most of these extant numerical models either use irregular wave forcing but simplify the hydrodynamics (as is the case for nearshore circulation models in the operational Navy), or allow more involved hydrodynamic formulations but reduce the forcing to that of monochromatic wave theory. Either modeling option can potentially lead to nearshore hydrodynamic predictions that do not exhibit the variability seen in nature.

Numerical Modeling of Nearshore Circulation: Recently, the Naval Research Laboratory (NRL) developed significant modifications to a sophisticated quasi-three-dimensional (quasi-3D) numerical hydrodynamic model. This model, SHORECIRC, was de-

veloped by Dr. Ib Svendsen and students at the Center for Applied Coastal Research of the University of Delaware. It is "quasi-3D" in that it actively models the depth-averaged horizontal velocities, but uses sophisticated analytic expressions to calculate the vertical profile of these horizontal velocities. The model is therefore able to simulate the evolution of horizontal velocities of the nearshore circulation field (including the depth-varying offshore flow, or "undertow") in all three dimensions (and time) without large computational effort. However, its physical mechanisms were geared toward monochromatic (single frequency) wave forcing, and thus are not applicable to field cases where waves of many frequencies and directions are present. NRL performed the reformulation of these physical mechanisms by adopting a probability distribution of waveheights inside the surf zone, and then integrating existing monochromatic formulations over this distribution. The resulting model represents the first implementation of random wave forcing in a general quasi-3D hydrodynamic model, thus allowing direct application to field situations.

Application to Nearshore Canyon: A major nearshore measurement campaign for the 2003-2004 time frame will be conducted near the Scripps Institution of Oceanography in La Jolla, California. This field experiment will be a collaborative effort among several universities and research institutions, including NRL. The site of the experiment is located at the head of Scripps Canyon, a major undersea canyon that exerts a strong polarizing effect on the nearshore wave and circulation climate. Figure 3 shows an aerial photograph of Black's Beach, located at the head of Scripps Canyon, taken by Dr. Steve Elgar of Woods Hole Oceanographic Institution. The



FIGURE 3

Aerial photograph of Black's Beach, north of Scripps Canyon in California. The undersea canyon enacts a strong variability on the nearshore wave and circulation environment. Arrows trace the presumed longshore and rip current patterns. Photo courtesy of Dr. Steve Elgar, Woods Hole Oceanographic Institution.

high variability in the wavefield and complex rip current environment is evident.

NRL developed a nested modeling system to investigate the possible nearshore flow patterns at this site, using the reformulated SHORECIRC model at the finest nest. The coastal-scale wave model SWAN (developed at Delft Technical University) propagated offshore wave energy into the nearshore canyon site. Figure 4 shows the waveheight prediction from SWAN over the entire domain, which was resolved at ~30 m in both directions. Information from SWAN was then input to a high-resolution nearshore irregular wave model (REF/DIF-S, developed at the University of Delaware) for simulating wave propagation over the small area in the white box near Black's Beach (at ~4-m resolution in both directions). The high-resolution wave model results were then used for forcing in the reformulated SHORECIRC model. Figure 5 shows the predicted average nearshore circulation pattern. The pair of northward-directed rip currents qualitatively represents what is seen in Fig.

3. No attempt was made to simulate the exact condition depicted in Fig. 3, so the rip current pair is probably more a direct result of the complex bathymetry than the wave climate offshore of the domain, and thus is likely to be a persistent feature.

Summary: NRL has developed significant modifications to the quasi-3D hydrodynamic model SHORECIRC. The most essential of these modifications (irregular wave forcing) enables the modeling of nearshore circulation without gross simplification of actual conditions. NRL developed a nested modeling system for the Scripps Canyon area, the site of an upcoming measurement campaign, and used this system to investigate potential flow conditions in the area. The presence of a rip current pair in the results qualitatively agrees with observations in the area. The NRL-enhanced SHORECIRC model shows promise as a forecasting tool, and further model validation is aimed toward this evaluation.

[Sponsored by ONR]




FIGURE 4

NRL model nest at Scripps Canyon: waveheights from the SWAN wave model over the shelf area near Scripps Canyon and Black's Beach. North is upward. Wave spectra with offshore height of 2 m and peak period of 17 s approach from the northwest. Waveheights over the domain range from 4 m (darkest red) to 0.25 m (deepest blue). Results are input into high-resolution wave and hydrodynamic models near Black's Beach (white rectangle).

FIGURE 5

NRL model nest at Scripps Canyon: predicted nearshore circulation pattern at Black's Beach from NRL-modified SHORECIRC model. Colors denote magnitude of longshore (north-south) velocities; range is from 1 m/s northward (dark red) to 1 m/s southward (deep blue). Arrows denote nearshore circulation pattern. Rip current pair (white arrows) are persistent features in simulations in this area, thus suggesting strong bathymetric influence on current patterns and milder sensitivity to nature of offshore conditions.

